

CORRIGENDUM

DALE R. DURRAN

Department of Atmospheric Sciences, University of Washington, Seattle, Washington

MARK GINGRICH

Apple Computer, Inc., Cupertino, California

(Manuscript received 8 December 2014, in final form 23 December 2014)

The quantitative comparison of initial errors in the velocity fields at 400 km, 10 km, and 10 cm in [Durrán and Gingrich \(2014\)](#) are incorrect because the square-root operator was not applied to convert kinetic energy (KE) values to velocities. The required quantitative corrections do not change our basic conclusions. The second paragraph of the conclusions should be revised as follows:

As evident from experiment B in L69 [[Lorenz \(1969\)](#)], but largely overlooked since, a small absolute error in the KE' spectral density produces almost the same loss in predictability no matter what its scale. Since the background saturation kinetic energy density is much bigger at longer wavelengths, very small relative errors in the large scales can have the same impact on predictability as saturated errors in the small scales. For example, consider a relative error of 100% in both the perturbation velocities and the KE' spectral density at a wavelength of 10 km. Assuming a $k^{-5/3}$ KE spectrum, the same absolute error in the perturbation velocities at 400 km corresponds to a relative error of $\{[(2\pi/10)/(2\pi/400)]^{-5/3}\}^{1/2}$. Thus, according to L69, RS08 [[Rotunno and Snyder \(2008\)](#)], and the smooth-saturation Lorenz–Rotunno–Snyder (ssLRS) models, 4.6% errors in velocities around nominal scales of 400 km would have a similar impact on predictability as 100% errors in velocities at scales around 10 km. If one pushes the comparison well past the limits of validity of the ssLRS model and imagines that butterflies all over the world are flapping in coordination to generate a 100% relative error at a wavelength of 10 cm, a roughly equivalent impact on predictability would be exerted by a tiny $3.1 \times 10^{-4}\%$ relative error in the perturbation velocities at a wavelength of 400 km. In any real-world event, the contributions of butterflies to uncertainties in initial conditions would be completely dwarfed by errors in the larger scales.

REFERENCES

- Durrán, D. R., and M. Gingrich, 2014: Atmospheric predictability: Why butterflies are not of practical importance. *J. Atmos. Sci.*, **71**, 2476–2488, doi:[10.1175/JAS-D-14-0007.1](#).
- Lorenz, E., 1969: The predictability of a flow which possesses many scales of motion. *Tellus*, **21**, 289–307, doi:[10.1111/j.2153-3490.1969.tb00444.x](#).
- Rotunno, R., and C. Snyder, 2008: A generalization of Lorenz's model for the predictability of flows with many scales of motion. *J. Atmos. Sci.*, **65**, 1063–1076, doi:[10.1175/2007JAS2449.1](#).

Corresponding author address: Dale Durrán, Dept. of Atmospheric Sciences, University of Washington, Box 351640, Seattle, WA 98195.
E-mail: drdee@uw.edu